

DATA DEPENDENCE RESULTS OF A NEW MULTISTEP AND S-ITERATIVE SCHEMES FOR CONTRACTIVE-LIKE OPERATORS

FAIK GURSOY, VATAN KARAKAYA, AND B. E. RHOADES

ABSTRACT. In this paper, we prove that convergence of a new iteration and S-iteration can be used approximate the fixed points of contractive-like operators. We also prove some data dependence results for these new iteration and S-iteration schemes for contractive-like operators. Our results extend and improve some known results in the literature.

1. INTRODUCTION

Contractive mappings and iteration procedures are some of the main tools in the study of fixed point theory. There are many contractive mappings and iteration schemes that have been introduced and developed by several authors to serve various purposes in the literature of this highly active research area, viz., [2, 39, 28, 17, 3, 33, 38, 23, 4, 40, 10, 27], among others.

Whether an iteration method used in any investigation converges to a fixed point of a contractive type mapping corresponding to a particular iteration process is of utmost importance. Therefore it is natural to see many works related to convergence of iteration methods, such as [37, 6, 7, 29, 9, 15, 24, 20, 1, 21].

Fixed point theory is concerned with investigating a wide variety of issues such as the existence (and uniqueness) of fixed points, the construction of fixed points, etc. One of these themes is data dependency of fixed points. Data dependency of fixed points has been the subject of research in fixed point theory for some time now, and data dependence research is an important theme in its own right.

Several authors who have made contributions to the study of data dependence of fixed points are Rus and Muresan [12], Rus et al. [13, 14], Berinde [36], Espínola and Petruşel [22], Markin [16], Chifu and Petruşel [5], Olantiwo [18, 19], Şoltuz [30, 31], Şoltuz and Grosan [32], Chugh and Kumar [25] and the references therein.

This paper is organized as follows. In Section 1 we present a brief survey of some known contractive mappings and iterative schemes and collect some preliminaries that will be used in the proofs of our main results. In Section 2 we show that the convergence of a new multi-step iteration, which is a special case of Jungck multistep-SP iterative process defined in [11] and S-iteration (due to Agarwal et al.) can be used approximate the fixed points of contractive-like operators. Motivated by the works of Şoltuz [30, 31], Şoltuz and Grosan [32], and Chugh and Kumar

Date: December, 2012.

2000 Mathematics Subject Classification. Primary 05C38, 15A15; Secondary 05A15, 15A18.

Key words and phrases. New multistep iteration, S-iteration, Data dependence, Contractive-like operator.

[25], we prove two data dependence results for the new multi-step iteration and S-iteration schemes by employing contractive-like operators.

As a background of our exposition, we now mention some contractive mappings and iteration schemes.

In [34] Zamfirescu established an important generalization of the Banach fixed point theorem using the following contractive condition: For a mapping $T : E \rightarrow E$, there exist real numbers a, b, c satisfying $0 < a < 1$, $0 < b, c < 1/2$ such that, for each pair $x, y \in X$, at least one of the following is true:

$$(1.1) \quad \begin{cases} (z_1) & \|Tx - Ty\| \leq a \|x - y\|, \\ (z_2) & \|Tx - Ty\| \leq b (\|x - Tx\| + \|y - Ty\|), \\ (z_3) & \|Tx - Ty\| \leq c (\|x - Ty\| + \|y - Tx\|). \end{cases}$$

A mapping T satisfying the contractive conditions (z_1) , (z_2) and (z_3) in (1.1) is called a Zamfirescu operator. An operator satisfying condition (z_2) is called a *Kannan operator*, while the mapping satisfying condition (z_3) is called a *Chatterjea operator*. As shown in [37], the contractive condition (1.1) leads to

$$(1.2) \quad \begin{cases} (b_1) & \|Tx - Ty\| \leq \delta \|x - y\| + 2\delta \|x - Tx\| \text{ if one use } (z_2), \\ \text{and} & \\ (b_2) & \|Tx - Ty\| \leq \delta \|x - y\| + 2\delta \|x - Ty\| \text{ if one use } (z_3), \end{cases}$$

for all $x, y \in E$ where $\delta := \max \left\{ a, \frac{b}{1-b}, \frac{c}{1-c} \right\}$, $\delta \in [0, 1)$, and it was shown that this class of operators is wider than the class of Zamfirescu operators. Any mapping satisfying condition (b_1) or (b_2) is called a quasi-contractive operator.

Extending the above definition, Osilike and Udomene [20] considered operators T for which there exist real numbers $L \geq 0$ and $\delta \in [0, 1)$ such that for all $x, y \in E$,

$$(1.3) \quad \|Tx - Ty\| \leq \delta \|x - y\| + L \|x - Tx\|.$$

Imoru and Olantiwo [8] gave a more general definition: The operator T is called a contractive-like operator if there exists a constant $\delta \in [0, 1)$ and a strictly increasing and continuous function $\varphi : [0, \infty) \rightarrow [0, \infty)$ with $\varphi(0) = 0$, such that, for each $x, y \in E$,

$$(1.4) \quad \|Tx - Ty\| \leq \delta \|x - y\| + \varphi(\|x - Tx\|).$$

A map satisfying (1.4) need not have a fixed point, even if E is complete. For example, let $E = [0, \infty)$, and define T by

$$Tx = \begin{cases} 1.0, & 0 \leq x \leq 0.8, \\ 0.6, & 0.8 < x. \end{cases}$$

Without loss of generality we may assume that $x < y$. Then, for $0 \leq x < y \leq 0.8$ or $0.8 < x < y$, $\|Tx - Ty\| = 0$, and (1.4) automatically satisfied.

If $0 \leq x \leq 0.8 < y$, then $\|Tx - Ty\| = 0.4$.

Define φ by $\varphi(t) = Lt$ for any $L \geq 2$. Then φ is increasing, continuous, and $\varphi(0) = 0$. Also, $\|x - Tx\| = 1 - x$, so that $\varphi(\|x - Tx\|) = L(1 - x) \geq 0.2L \geq 0.4$.

Therefore

$$0.4 = \|Tx - Ty\| \leq L \|x - Tx\| \leq \delta \|x - y\| + L \|x - Tx\|$$

for any $0 \leq \delta < 1$, and (1.4) is satisfied for $0 \leq x \leq 0.8 < y$. But T has no fixed point.

However, using (1.4) it is obvious that, if T has a fixed point, then it is unique.

Throughout this paper \mathbb{N} denotes the set of all nonnegative integers. Let X be a Banach space, $E \subset X$ a nonempty closed, convex subset of X , and T a self map on E . Define $F_T := \{p \in X : p = Tp\}$ to be the set of fixed points of T . Let $\{\alpha_n\}_{n=0}^\infty$, $\{\beta_n\}_{n=0}^\infty$, $\{\gamma_n\}_{n=0}^\infty$ and $\{\beta_n^i\}_{n=0}^\infty$, $i = \overline{1, k-2}$, $k \geq 2$ be real sequences in $[0, 1)$ satisfying certain conditions.

In [3] Rhoades and Soltuz introduced a multi-step iterative procedure defined by

$$(1.5) \quad \begin{cases} x_0 \in E, \\ y_n^{k-1} = (1 - \beta_n^{k-1})x_n + \beta_n^{k-1}Tx_n, \quad k \geq 2, \\ y_n^i = (1 - \beta_n^i)x_n + \beta_n^iTy_n^{i+1}, \quad i = \overline{1, k-2}, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_nTy_n^1, \quad n \in \mathbb{N}. \end{cases}$$

The sequence $\{x_n\}_{n=0}^\infty$ defined by

$$(1.6) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)Tx_n + \alpha_nTy_n, \\ y_n = (1 - \beta_n)x_n + \beta_nTx_n, \quad n \in \mathbb{N}, \end{cases}$$

is known as the S-iteration process (see [9, 26, 27]).

S.Thianwan [33] defined a two-step iteration $\{u_n\}_{n=0}^\infty$ by

$$(1.7) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)y_n + \alpha_nTy_n, \\ y_n = (1 - \beta_n)x_n + \beta_nTx_n, \quad n \in \mathbb{N}. \end{cases}$$

Recently Phuengrattana and Suantai [38] introduced an SP iteration method defined by

$$(1.8) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)y_n + \alpha_nTy_n, \\ y_n = (1 - \beta_n)z_n + \beta_nTz_n, \\ z_n = (1 - \gamma_n)x_n + \gamma_nTx_n, \quad n \in \mathbb{N}. \end{cases}$$

We shall employ the following iterative process. For an arbitrary fixed order $k \geq 2$,

$$(1.9) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)y_n^1 + \alpha_nTy_n^1, \\ y_n^1 = (1 - \beta_n^1)y_n^2 + \beta_n^1Ty_n^2, \\ y_n^2 = (1 - \beta_n^2)y_n^3 + \beta_n^2Ty_n^3, \\ \dots \\ y_n^{k-2} = (1 - \beta_n^{k-2})y_n^{k-1} + \beta_n^{k-2}Ty_n^{k-1}, \\ y_n^{k-1} = (1 - \beta_n^{k-1})x_n + \beta_n^{k-1}Tx_n, \quad n \in \mathbb{N}, \end{cases}$$

or, in short,

$$(1.10) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)y_n^1 + \alpha_nTy_n^1, \\ y_n^i = (1 - \beta_n^i)y_n^{i+1} + \beta_n^iTy_n^{i+1}, \quad i = \overline{1, k-2}, \\ y_n^{k-1} = (1 - \beta_n^{k-1})x_n + \beta_n^{k-1}Tx_n, \quad n \in \mathbb{N}, \end{cases}$$

where

$$(1.11) \quad \{\alpha_n\}_{n=0}^\infty \subset [0, 1), \quad \sum_{n=0}^\infty \alpha_n = \infty,$$

and

$$(1.12) \quad \{\beta_n^i\}_{n=0}^\infty \subset [0, 1), i = \overline{1, k-1}.$$

Remark 1. If $\gamma_n = 0$, then SP iteration (1.8) reduces to the two-step iteration (1.7). By taking $k = 3$ and $k = 2$ in (1.10) we obtain the iterations (1.8) and (1.7), respectively.

We shall need following definition and lemma in the sequel.

Definition 1. [35] Let $T, \tilde{T} : X \rightarrow X$ be two operators. We say that \tilde{T} is an approximate operator for T if, for some $\varepsilon > 0$, we have

$$\|Tx - \tilde{T}x\| \leq \varepsilon,$$

for all $x \in X$.

Lemma 1. [32] Let $\{a_n\}_{n=0}^\infty$ be a nonnegative sequence for which one assumes that there exists an $n_0 \in \mathbb{N}$, such that, for all $n \geq n_0$,

$$a_{n+1} \leq (1 - \mu_n) a_n + \mu_n \eta_n$$

is satisfied, where $\mu_n \in (0, 1)$, for all $n \in \mathbb{N}$, $\sum_{n=0}^\infty \mu_n = \infty$ and $\eta_n \geq 0$, $\forall n \in \mathbb{N}$.

Then the following holds:

$$0 \leq \limsup_{n \rightarrow \infty} a_n \leq \limsup_{n \rightarrow \infty} \eta_n.$$

2. MAIN RESULTS

For simplicity we use the following notation through this section.

For any iterative process, $\{x_n\}_{n=0}^\infty$ and $\{u_n\}_{n=0}^\infty$ denote iterative sequences associated to T and \tilde{T} , respectively.

Theorem 1. Let $T : E \rightarrow E$ be a map satisfying (1.4) with $F_T \neq \emptyset$ and $\{x_n\}_{n=0}^\infty$ is a sequence defined by (1.10), then the sequence $\{x_n\}_{n=0}^\infty$ converges to the unique fixed point of T .

Proof. The proof can be easily obtained by using argument in the proof of ([11], Theorem 3.1). \square

This result allow us to prove the following theorem.

Theorem 2. Let $T : E \rightarrow E$ be a map satisfying (1.4) with $F_T \neq \emptyset$ and \tilde{T} be an approximate operator of T as in the Definition 1. Let $\{x_n\}_{n=0}^\infty, \{u_n\}_{n=0}^\infty$ be two iterative sequences defined by (1.10) and with real sequences $\{\alpha_n\}_{n=0}^\infty, \{\beta_n^i\}_{n=0}^\infty \subset [0, 1)$ satisfying (i) $0 \leq \beta_n^i < \alpha_n \leq 1$, $i = \overline{1, k-1}$, (ii) $\sum \alpha_n = \infty$. If $p = Tp$ and $q = \tilde{T}q$, then we have

$$\|p - q\| \leq \frac{k\varepsilon}{1 - \delta}.$$

Proof. For a given $x_0 \in E$ and $u_0 \in E$ we consider following multistep iteration for T and \tilde{T} :

$$(2.1) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n) y_n^1 + \alpha_n T y_n^1, \\ y_n^i = (1 - \beta_n^i) y_n^{i+1} + \beta_n^i T y_n^{i+1}, i = \overline{1, k-2}, \\ y_n^{k-1} = (1 - \beta_n^{k-1}) x_n + \beta_n^{k-1} T x_n, k \geq 2, n \in \mathbb{N}, \end{cases}$$

and

$$(2.2) \quad \begin{cases} u_0 \in E, \\ u_{n+1} = (1 - \alpha_n) v_n^1 + \alpha_n \tilde{T} v_n^1, \\ v_n^i = (1 - \beta_n^i) v_n^{i+1} + \beta_n^i \tilde{T} v_n^{i+1}, \quad i = \overline{1, k-2}, \\ v_n^{k-1} = (1 - \beta_n^{k-1}) u_n + \beta_n^{k-1} \tilde{T} u_n, \quad k \geq 2, \quad n \in \mathbb{N}. \end{cases}$$

Then, from (1.4), (2.1) and (2.2), we have the following estimates.

$$(2.3) \quad \begin{aligned} \|x_{n+1} - u_{n+1}\| &= \left\| (1 - \alpha_n) (y_n^1 - v_n^1) + \alpha_n (Ty_n^1 - \tilde{T}v_n^1) \right\| \\ &\leq (1 - \alpha_n) \|y_n^1 - v_n^1\| + \alpha_n \|Ty_n^1 - \tilde{T}v_n^1\| \\ &= (1 - \alpha_n) \|y_n^1 - v_n^1\| + \alpha_n \|Ty_n^1 - Tv_n^1 + Tv_n^1 - \tilde{T}v_n^1\| \\ &\leq (1 - \alpha_n) \|y_n^1 - v_n^1\| + \alpha_n \|Ty_n^1 - Tv_n^1\| + \alpha_n \|Tv_n^1 - \tilde{T}v_n^1\| \\ &\leq (1 - \alpha_n) \|y_n^1 - v_n^1\| + \alpha_n \delta \|y_n^1 - v_n^1\| + \alpha_n \varphi(\|y_n^1 - Ty_n^1\|) + \alpha_n \varepsilon \\ &= [1 - \alpha_n (1 - \delta)] \|y_n^1 - v_n^1\| + \alpha_n \varphi(\|y_n^1 - Ty_n^1\|) + \alpha_n \varepsilon, \end{aligned}$$

$$(2.4) \quad \begin{aligned} \|y_n^1 - v_n^1\| &= \left\| (1 - \beta_n^1) (y_n^2 - v_n^2) + \beta_n^1 (Ty_n^2 - \tilde{T}v_n^2) \right\| \\ &\leq (1 - \beta_n^1) \|y_n^2 - v_n^2\| + \beta_n^1 \|Ty_n^2 - \tilde{T}v_n^2\| \\ &\leq (1 - \beta_n^1) \|y_n^2 - v_n^2\| + \beta_n^1 \|Ty_n^2 - Tv_n^2\| + \beta_n^1 \|Tv_n^2 - \tilde{T}v_n^2\| \\ &\leq (1 - \beta_n^1) \|y_n^2 - v_n^2\| + \beta_n^1 \delta \|y_n^2 - v_n^2\| + \beta_n^1 \varphi(\|y_n^2 - Ty_n^2\|) + \beta_n^1 \varepsilon \\ &= [1 - \beta_n^1 (1 - \delta)] \|y_n^2 - v_n^2\| + \beta_n^1 \varphi(\|y_n^2 - Ty_n^2\|) + \beta_n^1 \varepsilon, \end{aligned}$$

$$(2.5) \quad \begin{aligned} \|y_n^2 - v_n^2\| &= \left\| (1 - \beta_n^2) (y_n^3 - v_n^3) + \beta_n^2 (Ty_n^3 - \tilde{T}v_n^3) \right\| \\ &\leq (1 - \beta_n^2) \|y_n^3 - v_n^3\| + \beta_n^2 \|Ty_n^3 - \tilde{T}v_n^3\| \\ &\leq (1 - \beta_n^2) \|y_n^3 - v_n^3\| + \beta_n^2 \|Ty_n^3 - Tv_n^3\| + \beta_n^2 \|Tv_n^3 - \tilde{T}v_n^3\| \\ &\leq (1 - \beta_n^2) \|y_n^3 - v_n^3\| + \beta_n^2 \delta \|y_n^3 - v_n^3\| + \beta_n^2 \varphi(\|y_n^3 - Ty_n^3\|) + \beta_n^2 \varepsilon \\ &= [1 - \beta_n^2 (1 - \delta)] \|y_n^3 - v_n^3\| + \beta_n^2 \varphi(\|y_n^3 - Ty_n^3\|) + \beta_n^2 \varepsilon. \end{aligned}$$

Combining (2.3), (2.4) and (2.5) we obtain

$$(2.6) \quad \begin{aligned} \|x_{n+1} - u_{n+1}\| &\leq [1 - \alpha_n (1 - \delta)] [1 - \beta_n^1 (1 - \delta)] [1 - \beta_n^2 (1 - \delta)] \|y_n^3 - v_n^3\| \\ &\quad + [1 - \alpha_n (1 - \delta)] [1 - \beta_n^1 (1 - \delta)] \beta_n^2 \varphi(\|y_n^3 - Ty_n^3\|) \\ &\quad + [1 - \alpha_n (1 - \delta)] [1 - \beta_n^1 (1 - \delta)] \beta_n^2 \varepsilon \\ &\quad + [1 - \alpha_n (1 - \delta)] \beta_n^1 \varphi(\|y_n^2 - Ty_n^2\|) \\ &\quad + [1 - \alpha_n (1 - \delta)] \beta_n^1 \varepsilon + \alpha_n \varphi(\|y_n^1 - Ty_n^1\|) + \alpha_n \varepsilon. \end{aligned}$$

Thus, by induction, we get

$$\begin{aligned}
\|x_{n+1} - u_{n+1}\| &\leq [1 - \alpha_n (1 - \delta)] \\
&\quad [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-2} (1 - \delta)] \|y_n^{k-1} - v_n^{k-1}\| \\
&\quad + [1 - \alpha_n (1 - \delta)] \\
&\quad [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-3} (1 - \delta)] \beta_n^{k-2} \varphi (\|y_n^{k-1} - Ty_n^{k-1}\|) \\
&\quad + \cdots + [1 - \alpha_n (1 - \delta)] \beta_n^1 \varphi (\|y_n^2 - Ty_n^2\|) + \alpha_n \varphi (\|y_n^1 - Ty_n^1\|) \\
&\quad + [1 - \alpha_n (1 - \delta)] [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-3} (1 - \delta)] \beta_n^{k-2} \varepsilon \\
(2.7) \quad &\quad + \cdots + [1 - \alpha_n (1 - \delta)] \beta_n^1 \varepsilon + \alpha_n \varepsilon.
\end{aligned}$$

Again using (1.4), (2.1) and (2.2) we get

$$\begin{aligned}
\|y_n^{k-1} - v_n^{k-1}\| &= \left\| (1 - \beta_n^{k-1}) (x_n - u_n) + \beta_n^{k-1} (Tx_n - \tilde{T}u_n) \right\| \\
&\leq (1 - \beta_n^{k-1}) \|x_n - u_n\| + \beta_n^{k-1} \|Tx_n - \tilde{T}u_n\| \\
&\leq (1 - \beta_n^{k-1}) \|x_n - u_n\| + \beta_n^{k-1} \|Tx_n - Tu_n\| \\
&\quad + \beta_n^{k-1} \|Tu_n - \tilde{T}u_n\| \\
(2.8) \quad &\leq [1 - \beta_n^{k-1} (1 - \delta)] \|x_n - u_n\| + \beta_n^{k-1} \varphi (\|x_n - Tx_n\|) + \beta_n^{k-1} \varepsilon.
\end{aligned}$$

Substituting (2.8) in (2.7) we have

$$\begin{aligned}
\|x_{n+1} - u_{n+1}\| &\leq [1 - \alpha_n (1 - \delta)] \\
&\quad [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-1} (1 - \delta)] \|x_n - u_n\| \\
&\quad + [1 - \alpha_n (1 - \delta)] \\
&\quad [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-2} (1 - \delta)] \beta_n^{k-1} \varphi (\|x_n - Tx_n\|) \\
&\quad + [1 - \alpha_n (1 - \delta)] \\
&\quad [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-3} (1 - \delta)] \beta_n^{k-2} \varphi (\|y_n^{k-1} - Ty_n^{k-1}\|) \\
&\quad + \cdots + [1 - \alpha_n (1 - \delta)] \beta_n^1 \varphi (\|y_n^2 - Ty_n^2\|) + \alpha_n \varphi (\|y_n^1 - Ty_n^1\|) \\
&\quad + [1 - \alpha_n (1 - \delta)] [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-2} (1 - \delta)] \beta_n^{k-1} \varepsilon \\
(2.9) \quad &\quad + \cdots + [1 - \alpha_n (1 - \delta)] \beta_n^1 \varepsilon + \alpha_n \varepsilon.
\end{aligned}$$

Since $\delta \in [0, 1)$ and $\{\alpha_n\}_{n=0}^\infty, \{\beta_n^i\}_{n=0}^\infty \subset [0, 1)$ for $i = \overline{1, k-1}$, we have

$$(2.10) \quad [1 - \alpha_n (1 - \delta)] [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^i (1 - \delta)] \leq [1 - \alpha_n (1 - \delta)].$$

Using inequality (2.10) and assumption (i) in (2.9) it follows

$$\begin{aligned}
 \|x_{n+1} - u_{n+1}\| &\leq [1 - \alpha_n(1 - \delta)] \|x_n - u_n\| \\
 &\quad + \alpha_n \varphi(\|x_n - Tx_n\|) + \alpha_n \varphi(\|y_n^{k-1} - Ty_n^{k-1}\|) \\
 &\quad + \cdots + \alpha_n \varphi(\|y_n^2 - Ty_n^2\|) + \alpha_n \varphi(\|y_n^1 - Ty_n^1\|) \\
 &\quad + \alpha_n \varepsilon + \alpha_n \varepsilon + \cdots + \alpha_n \varepsilon + \alpha_n \varepsilon \\
 &= [1 - \alpha_n(1 - \delta)] \|x_n - u_n\| \\
 &\quad + \alpha_n(1 - \delta) \left\{ \frac{\varphi(\|x_n - Tx_n\|) + \varphi(\|y_n^{k-1} - Ty_n^{k-1}\|)}{1 - \delta} \right. \\
 &\quad \left. + \cdots + \frac{\varphi(\|y_n^1 - Ty_n^1\|) + k\varepsilon}{1 - \delta} \right\}.
 \end{aligned}
 \tag{2.11}$$

Define

$$\begin{aligned}
 a_n &: = \|x_n - u_n\|, \\
 \mu_n &: = \alpha_n(1 - \delta) \in (0, 1), \\
 \eta_n &: = \left\{ \frac{\varphi(\|x_n - Tx_n\|) + \varphi(\|y_n^{k-1} - Ty_n^{k-1}\|)}{1 - \delta} \right. \\
 &\quad \left. + \cdots + \frac{\varphi(\|y_n^1 - Ty_n^1\|) + k\varepsilon}{1 - \delta} \right\}.
 \end{aligned}$$

From Theorem 1 it follows that $\lim_{n \rightarrow \infty} \|x_n - p\| = 0$. Since T satisfies condition (1.4) and $Tp = p \in F_T$,

$$\begin{aligned}
 0 &\leq \|x_n - Tx_n\| \\
 &\leq \|x_n - p\| + \|Tp - Tx_n\| \\
 &\leq \|x_n - p\| + \delta \|p - x_n\| + \varphi(\|p - Tp\|) \\
 &= (1 + \delta) \|x_n - p\| \rightarrow 0 \text{ as } n \rightarrow \infty.
 \end{aligned}
 \tag{2.12}$$

Since $\beta_n^i \in [0, 1]$, $\forall n \in \mathbb{N}$, $i = \overline{1, k-1}$ and using (1.4) and (1.10) we have

$$\begin{aligned}
 0 &\leq \|y_n^1 - Ty_n^1\| = \|y_n^1 - p + p - Ty_n^1\| \\
 &\leq \|y_n^1 - p\| + \|Tp - Ty_n^1\| \\
 &\leq \|y_n^1 - p\| + \delta \|p - y_n^1\| + \varphi(\|p - Tp\|) \\
 &= (1 + \delta) \|y_n^1 - p\| \\
 &= (1 + \delta) \|(1 - \beta_n^1) y_n^2 + \beta_n^1 Ty_n^2 - p(1 - \beta_n^1 + \beta_n^1)\| \\
 &\leq (1 + \delta) \{(1 - \beta_n^1) \|y_n^2 - p\| + \beta_n^1 \|Ty_n^2 - Tp\|\} \\
 &\leq (1 + \delta) \{(1 - \beta_n^1) \|y_n^2 - p\| + \beta_n^1 \delta \|y_n^2 - p\|\} \\
 &= (1 + \delta) [1 - \beta_n^1(1 - \delta)] \|y_n^2 - p\| \\
 &\leq \cdots \\
 &\leq (1 + \delta) [1 - \beta_n^1(1 - \delta)] \cdots [1 - \beta_n^{k-2}(1 - \delta)] \|y_n^{k-1} - p\| \\
 &\leq (1 + \delta) [1 - \beta_n^1(1 - \delta)] \cdots [1 - \beta_n^{k-1}(1 - \delta)] \|x_n - p\| \\
 &\leq (1 + \delta) \|x_n - p\| \rightarrow 0 \text{ as } n \rightarrow \infty.
 \end{aligned}
 \tag{2.13}$$

It is easy to see from (2.13) that this result is also valid for $\|Ty_n^2 - y_n^2\|, \dots, \|Ty_n^{k-1} - y_n^{k-1}\|$. Since φ is continuous, we have

$$(2.14) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \varphi(\|x_n - Tx_n\|) \\ &= \lim_{n \rightarrow \infty} \varphi(\|y_n^1 - Ty_n^1\|) = \dots = \lim_{n \rightarrow \infty} \varphi(\|y_n^{k-1} - Ty_n^{k-1}\|) = 0. \end{aligned}$$

Hence an application of Lemma 1 to (2.11) leads to

$$(2.15) \quad \|p - q\| \leq \frac{k\varepsilon}{1 - \delta}.$$

□

As shown by Hussain et al. ([21], Theorem 8), in an arbitrary Banach space X , the S-iteration $\{x_n\}_{n=0}^\infty$ given by (1.6) converges to the fixed point of T , where $T : E \rightarrow E$ is a mapping satisfying condition (1.3).

Theorem 3. *Let $T : E \rightarrow E$ be a map satisfying (1.4) with $F_T \neq \emptyset$ and $\{x_n\}_{n=0}^\infty$ be defined by (1.6) with real sequences $\{\beta_n\}_{n=0}^\infty$, $\{\alpha_n\}_{n=0}^\infty \subset [0, 1)$ satisfying $\sum_{n=0}^\infty \alpha_n = \infty$. Then the sequence $\{x_n\}_{n=0}^\infty$ converges to the unique fixed point of T .*

Proof. The argument is similar to the proof of Theorem 8 of [21], and is thus omitted. □

We now prove result on data dependence for the S-iterative procedure by utilizing Theorem 3.

Theorem 4. *Let T, \tilde{T} be two operators as in Theorem 2. Let $\{x_n\}_{n=0}^\infty, \{u_n\}_{n=0}^\infty$ be S-iterations defined by (1.6) and with real sequences $\{\beta_n\}_{n=0}^\infty, \{\alpha_n\}_{n=0}^\infty \subset [0, 1)$ satisfying (i) $\frac{1}{2} \leq \alpha_n, \forall n \in \mathbb{N}$, and (ii) $\sum_{n=0}^\infty \alpha_n = \infty$. If $p = Tp$ and $q = \tilde{T}q$, then we have*

$$\|p - q\| \leq \frac{3\varepsilon}{1 - \delta}.$$

Proof. For a given $x_0 \in C$ and $u_0 \in C$ we consider following iteration for T and \tilde{T} :

$$(2.16) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)Tx_n + \alpha_nTy_n, \\ y_n = (1 - \beta_n)x_n + \beta_nTx_n, n \in \mathbb{N} \end{cases}$$

and

$$(2.17) \quad \begin{cases} x_0 \in E, \\ u_{n+1} = (1 - \alpha_n)\tilde{T}u_n + \alpha_n\tilde{T}v_n, \\ v_n = (1 - \beta_n)u_n + \beta_n\tilde{T}u_n, n \in \mathbb{N}. \end{cases}$$

Using (1.4), (2.16) and (2.17), we obtain the following estimates

$$\begin{aligned}
\|x_{n+1} - u_{n+1}\| &= \left\| (1 - \alpha_n) (Tx_n - \tilde{T}u_n) + \alpha_n (Ty_n - \tilde{T}v_n) \right\| \\
&\leq (1 - \alpha_n) \|Tx_n - \tilde{T}u_n\| + \alpha_n \|Ty_n - \tilde{T}v_n\| \\
&= (1 - \alpha_n) \|Tx_n - Tu_n + Tu_n - \tilde{T}u_n\| \\
&\quad + \alpha_n \|Ty_n - Tv_n + Tv_n - \tilde{T}v_n\| \\
&\leq (1 - \alpha_n) \left\{ \|Tx_n - Tu_n\| + \|Tu_n - \tilde{T}u_n\| \right\} \\
&\quad + \alpha_n \left\{ \|Ty_n - Tv_n\| + \|Tv_n - \tilde{T}v_n\| \right\} \\
&\leq (1 - \alpha_n) \{ \delta \|x_n - u_n\| + \varphi(\|x_n - Tx_n\|) + \varepsilon \} \\
&\quad + \alpha_n \{ \delta \|y_n - v_n\| + \varphi(\|y_n - Ty_n\|) + \varepsilon \},
\end{aligned} \tag{2.18}$$

$$\begin{aligned}
\|y_n - v_n\| &= \left\| (1 - \beta_n) (x_n - u_n) + \beta_n (Tx_n - \tilde{T}u_n) \right\| \\
&\leq (1 - \beta_n) \|x_n - u_n\| + \beta_n \|Tx_n - \tilde{T}u_n\| \\
&= (1 - \beta_n) \|x_n - u_n\| + \beta_n \|Tx_n - Tu_n + Tu_n - \tilde{T}u_n\| \\
&\leq (1 - \beta_n) \|x_n - u_n\| + \beta_n \left\{ \|Tx_n - Tu_n\| + \|Tu_n - \tilde{T}u_n\| \right\} \\
&\leq (1 - \beta_n) \|x_n - u_n\| + \beta_n \{ \delta \|x_n - u_n\| + \varphi(\|x_n - Tx_n\|) + \varepsilon \} \\
&= [1 - \beta_n(1 - \delta)] \|x_n - u_n\| + \beta_n \varphi(\|x_n - Tx_n\|) + \beta_n \varepsilon.
\end{aligned} \tag{2.19}$$

Combining (2.18) and (2.19),

$$\begin{aligned}
\|x_{n+1} - u_{n+1}\| &\leq \{ (1 - \alpha_n) \delta + \alpha_n \delta [1 - \beta_n(1 - \delta)] \} \|x_n - u_n\| \\
&\quad + \{ 1 - \alpha_n + \alpha_n \delta \beta_n \} \varphi(\|x_n - Tx_n\|) + \alpha_n \varphi(\|y_n - Ty_n\|) \\
&\quad + \alpha_n \delta \beta_n \varepsilon + (1 - \alpha_n) \varepsilon + \alpha_n \varepsilon.
\end{aligned} \tag{2.20}$$

For $\{\alpha_n\}_{n=0}^\infty$, $\{\beta_n\}_{n=0}^\infty \subset [0, 1)$ and $\delta \in [0, 1)$

$$(1 - \alpha_n) \delta < 1 - \alpha_n, \quad 1 - \beta_n(1 - \delta) < 1, \quad \alpha_n \delta \beta_n < \alpha_n. \tag{2.21}$$

It follows from assumption (i) that

$$1 - \alpha_n < \alpha_n, \forall n \in \mathbb{N}. \tag{2.22}$$

Therefore, combining (2.22) and (2.21) to (2.20) gives

$$\begin{aligned}
\|x_{n+1} - u_{n+1}\| &\leq [1 - \alpha_n(1 - \delta)] \|x_n - u_n\| \\
&\quad + 2\alpha_n \varphi(\|x_n - Tx_n\|) + \alpha_n \varphi(\|y_n - Ty_n\|) \\
&\quad + \alpha_n \varepsilon + \alpha_n \varepsilon + \alpha_n \varepsilon,
\end{aligned} \tag{2.23}$$

or, equivalently,

$$\begin{aligned}
\|x_{n+1} - u_{n+1}\| &\leq [1 - \alpha_n(1 - \delta)] \|x_n - u_n\| \\
&\quad + \alpha_n(1 - \delta) \frac{\{2\varphi(\|x_n - Tx_n\|) + \varphi(\|y_n - Ty_n\|) + 3\varepsilon\}}{1 - \delta}.
\end{aligned} \tag{2.24}$$

Now define

$$\begin{aligned} a_n &: = \|x_n - u_n\|, \\ \eta_n &: = \alpha_n (1 - \delta) \in (0, 1) \\ \rho_n &: = \frac{2\varphi(\|x_n - Tx_n\|) + \varphi(\|y_n - Ty_n\|) + 3\varepsilon}{1 - \delta}. \end{aligned}$$

From Theorem 3, we have $\lim_{n \rightarrow \infty} \|x_n - p\| = 0$. Since T satisfies condition (1.4), and $Tp = p \in F_T$, using an argument similar to that in the proof of Theorem 2

$$(2.25) \quad \lim_{n \rightarrow \infty} \|x_n - Tx_n\| = \lim_{n \rightarrow \infty} \|y_n - Ty_n\| = 0.$$

Using the fact that φ is continuous we have

$$(2.26) \quad \lim_{n \rightarrow \infty} \varphi(\|x_n - Tx_n\|) = \lim_{n \rightarrow \infty} \varphi(\|y_n - Ty_n\|) = 0.$$

An application of Lemma 1 to (2.24) leads to

$$(2.27) \quad \|p - q\| \leq \frac{3\varepsilon}{1 - \delta}.$$

□

3. CONCLUSION

Since the iterative schemes (1.7) and (1.8) are special cases the iterative process (1.10), Theorem 1 generalizes Theorem 2.1 of [24], and Theorem 2.1 of [15]. By taking $k = 3$ and $k = 2$ in Theorem 2, data dependence results for the iterative schemes (1.7) and (1.8) can be easily obtained. For $k = 3$, Theorem 2 reduces to Theorem 3.2 of [25]. Since condition (1.4) is more general than condition (1.3), Theorem 3 generalizes Theorem 8 of [21].

Acknowledgement 1. *This work is supported by Yıldız Technical University Scientific Research Projects Coordination Unit under project number BAPK 2012-07-03-DOP02.*

REFERENCES

- [1] A. Rafiq, "On the convergence of the three step iteration process in the class of quasi-contractive operators", Acta Mathematica Academiae Paedagogicae Nyiregyhaziensis, **22**(2006), 305-309.
- [2] B.E. Rhoades, "A comparison of various definitions of contractive mappings", Transactions of the American Mathematical Society, **226**(1977), 257-290.
- [3] B.E. Rhoades, S.M. Şoltuz, "The equivalence between Mann-Ishikawa iterations and multi-step iteration", Nonlinear Analysis, **58**(2004), 219-228.
- [4] B. Xu, M.A. Noor, "Ishikawa and Mann iteration process with errors for nonlinear strongly accretive operator equations", J. Math. Anal. Appl., **224**(1998), 91-101.
- [5] C. Chifu, G. Petruşel, "Existence and Data Dependence of Fixed Points and Strict Fixed Points for Contractive-Type Multivalued Operators", Fixed Point Theory and Applications, **2007**(2007), Article ID 034248, 8 pages.
- [6] C. E. Chidume, C. O. Chidume, "Convergence theorem for fixed points of uniformly continuous generalized phi-hemicontractive mappings", J. Math. Anal. Appl., **303**(2005), 545-554.
- [7] C.E. Chidume, C.O. Chidume, "Iterative approximation of fixed points of nonexpansive mappings", J. Math. Anal. Appl., **318**(2006), no. 1, 288-295.
- [8] C.O. Imoru, M.O. Olantiwo, "On the stability of Picard and Mann iteration processes", Carpathian Journal of Mathematics, **19**(2003), no.2, 155-160.
- [9] D.R. Sahu, "Applications of the S-iteration process to constrained minimization problems and split feasibility problems", Fixed Point Theory, **12**(2011), no. 1, 187-204.

- [10] G. Das, J.P. Debata, "Fixed points of Quasi-nonexpansive mappings", Indian J. Pure, Appl. Math., **17**(1986), 1263-1269.
- [11] H. Akewe, "Strong convergence and stability of Jungck-multistep-SP iteration for generalized contractive-like inequality operators", Advances in Natural Science, **5**(2012), no. 3, 21-27.
- [12] I.A. Rus, S. Muresan, "Data dependence of the fixed points set of weakly Picard operators", Stud. Univ. Babes-Bolyai **43** (1998) 79-83.
- [13] I.A. Rus, A. Petruşel, A. Sintamarian, "Data dependence of the fixed points set of multivalued weakly Picard operators", Stud. Univ. Babes-Bolyai, Math. **46** (2) (2001) 111-121.
- [14] I.A. Rus, A. Petruşel, A. Sintamarian, "Data dependence of the fixed point set of some multivalued weakly Picard operators", Nonlinear Analysis: Theory, Methods & Applications, **52** (2003), 1947-1959.
- [15] İ. Yıldırım, M. Özdemir, H. Kızıltuğ, "On the convergence of a new two-step iteration in the class of quasi-contractive operators", Int. Journal of Math. Analysis, **3**(2009), no. 38, 1881-1892.
- [16] J.T. Markin, "Continuous dependence of fixed point sets", Proc. AMS **38** (1973) 545-547.
- [17] M.A. Noor, "New approximation schemes for general variational inequalities", Journal of Mathematical Analysis and Applications, **251**(2000), no.1, 217-229.
- [18] M. O. Olatinwo, "Some results on the continuous dependence of the fixed points in normed linear space", Fixed Point Theory, **10**(2009), no. 1, 151-157.
- [19] M. O. Olatinwo, "On the continuous dependence of the fixed points for (φ, ψ) -contractive-type operators", Kragujevac Journal of Mathematics, **34**(2010), 91-102.
- [20] M.O. Osilike, A. Udomene, "Short proofs of stability results for fixed point iteration procedures for a class of contractive-type mappings", Indian Journal of Pure and Applied Mathematics, **30**(1999), no.12, 1229-1234.
- [21] N. Hussain, A. Rafiq, B. Damjanović, R. Lazović, "On rate of convergence of various iterative schemes", Fixed Point Theory and Applications, **2011**(2011), doi:10.1186/1687-1812-2011-45.
- [22] R. Espínola, A. Petruşel, "Existence and data dependence of fixed points for multivalued operators on gauge spaces", J. Math. Anal. Appl., **309** (2005), 420-432.
- [23] R. Glowinski, P. Le Tallec, "Augmented Lagrangian and operator splitting methods in non-linear mechanics", SIAM, Philadelphia, 1989.
- [24] R. Chugh, V. Kumar, "Strong convergence of SP iterative scheme for quasi-contractive operators in Banach spaces", International Journal of Computer Applications, **31**(2011), no.5.
- [25] R. Chugh, V. Kumar, "Data dependence of Noor and SP iterative schemes when dealing with quasi-contractive operators", International Journal of Computer Applications, **31**(2011), no.5.
- [26] R.P. Agarwal, D. O'Regan, D.R. Sahu, "Fixed point theory for lipschitzian type-mappings with applications", Springer, ISBN: 978-0-387-75817-6, 2009.
- [27] R.P. Agarwal, D. O'Regan, D.R. Sahu, "Iterative construction of fixed points of nearly asymptotically nonexpansive mappings", J. Nonlinear Convex Anal., **8**(2007), no.1, 61-79.
- [28] S. Ishikawa, "Fixed points by a new iteration method", Proc. Amer. Math. Soc., **44**(1974), 147-150.
- [29] S. Suantai, "Weak and strong convergence criteria of Noor iterations for asymptotically nonexpansive mappings", J. Math. Anal. Appl., **311**(2005), no.2, 506-517.
- [30] S.M. Şoltuz, "Data dependence for Mann iteration", Octogon Math. Magazine, **9**(2001), 825-828.
- [31] S.M. Şoltuz, "Data dependence for Ishikawa iteration", Lecturas Mathematicas, **25**(2004), no. 2, 149-155.
- [32] S.M. Şoltuz, T. Grosan, "Data dependence for Ishikawa iteration when dealing with contractive like operators", Fixed Point Theory and Applications, **2008**(2008), Article ID 242916, 7 pages.
- [33] S. Thianwan, "Common fixed points of new iterations for two asymptotically nonexpansive nonself mappings in a Banach space", J. Comput. Appl. Math., (2008), doi: 10.1016/j.cam.2008.05.051.
- [34] T. Zamfirescu, "Fix point theorems in metric spaces", Archiv der Mathematik, **23**(1972), no.1, 292-298.
- [35] V. Berinde, "Iterative Approximation of Fixed Points ", Springer, Berlin (2007).

- [36] V. Berinde, "*On the approximation of fixed points of weak contractive mappings*", Carpathian J. Math., **19**(2003), no. 1, 7-22.
- [37] V. Berinde, "*On the convergence of the Ishikawa iteration in the class of quasi contractive operators*", Acta Mathematica Universitatis Comenianae, **73**(2004), no.1, 119-126.
- [38] W. Phuengrattana, S. Suantai, "*On the rate of convergence of Mann, Ishikawa, Noor and SP iterations for continuous functions on an arbitrary interval*", Journal of Computational and Applied Mathematics, **235**(2011), 3006-3014.
- [39] W.R. Mann, "*Mean value methods in iterations*", Proc. Amer. Math. Soc., **4**(1953), 506-510.
- [40] W. Takahashi, "*Iterative methods for approximation of fixed points and their applications*", Journal of the Operations Research Society of Japan, **43**(2000), no.1, 87-108.

DEPARTMENT OF MATHEMATICS, YILDIZ TECHNICAL UNIVERSITY, DAVUTPASA CAMPUS, ESENLER, 34220 ISTANBUL, TURKEY

E-mail address: faikgursoy02@hotmail.com; fgursoy@yildiz.edu.tr

URL: <http://www.yarbis.yildiz.edu.tr/fgursoy>

Current address: Department of Mathematical Engineering, Yildiz Technical University, Davutpasa Campus, Esenler, 34210 Istanbul

E-mail address: vkkaya@yildiz.edu.tr; vkkaya@yahoo.com

URL: <http://www.yarbis.yildiz.edu.tr/vkkaya>

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405-7106, USA

E-mail address: rhoades@indiana.edu

URL: <http://www.math.indiana.edu/people/profile.phtml?id=rhoades>